

Coexistence of solutions in dynamical mean-field theory of the Mott transition

Werner Krauth *

CNRS-Laboratoire de Physique Statistique

Ecole Normale Supérieure, 24, rue Lhomond, 75231 Paris Cedex 05, France

In this paper, I discuss the finite-temperature metal-insulator transition of the paramagnetic Hubbard model within dynamical mean-field theory. I show that coexisting solutions, the hallmark of such a transition, can be obtained in a consistent way both from Quantum Monte Carlo (QMC) simulations and from the Exact Diagonalization method. I pay special attention to discretization errors within QMC. These errors explain why it is difficult to obtain the solutions by QMC close to the boundaries of the coexistence region.

PACS numbers: 71.27.+a

The Mott-Hubbard metal-insulator transition is one of the fundamental problems in the field of correlated electronic systems. During the last few years, dynamical mean-field theory (DMFT) [1] [2] has emerged as an appropriate paradigm for this transition. Within DMFT, the paramagnetic half-filled Hubbard model on the Bethe lattice was found to undergo a first-order phase transition at finite temperature $T = 1/\beta$. The hallmark of such a transition is the coexistence of two solutions for the same value of the electronic interaction U with $U_{c_1}(\beta) < U < U_{c_2}(\beta)$. As is usual, I will label “insulating” the solution which can be continuously followed at low temperature from large values of the interaction down to U_{c_1} and “metallic” the one which exists from $U = 0$ up to U_{c_2} .

Recent work [3] has challenged the above scenario. Using the Quantum Monte Carlo (QMC) method, rather than the Exact Diagonalization (ED) approach [4], the authors of [3] were unable to detect coexistence in the DMFT equations. This is surprising, as the two algorithms were compared extensively to each other (cf, e.g. ref. [2], section VI, Appendix C). The aim of the present paper is to discuss the conflicting numerical approaches (for related work, cf [5]). I show that coexisting metallic and insulating solutions can indeed be obtained using both methods in a transparent and completely consistent way. I pay special attention to the problem of discretization errors. It will become clear that the QMC method needs to use very fine discretizations in order to obtain the insulating solution close to U_{c_1} . Inside the coexistence region, the problem disappears. All the numerical work in this paper relies on programs which are publicly available [2].

Given the large number of articles already published on the subject, I will not repeat the standard definitions for the half-filled Hubbard model on the Bethe lattice within DMFT. These can be found e.g. in ref. [2]. The important parameter of the model is the bandwidth D (cf. [2]); I quote interactions and temperatures in units

of the bandwidth $D/\sqrt{2}$.

The point of departure of the present paper is ref. [6], where coexisting solutions of the mean-field equations at inverse temperature $\beta D/\sqrt{2} = 100$ were found (within ED) for $U_{c_1} < U < U_{c_2}$ with $U_{c_1}(\beta D/\sqrt{2} = 100) \sim 3.3D/\sqrt{2}$ and $U_{c_2}(\beta D/\sqrt{2} = 100) = 3.8D/\sqrt{2}$.

For this paper, I choose as a reference point the value $U = 3.55D/\sqrt{2}$, $\beta D/\sqrt{2} = 100$. At this point, both metallic and insulating solutions can be found easily by ED [7]. These solutions are given by bath Green’s functions $\mathcal{G}_0(\tau)$ and impurity model Green’s functions $G(\tau)$ (cf [2] for definitions).

As we are well within the coexistence region, both solutions correspond to the bottoms of deep basins of attraction. I have, e.g., perturbed both the metallic bath Green’s function, $\mathcal{G}_0^{met}(\tau)$, as well as the insulating one, $\mathcal{G}_0^{ins}(\tau)$, and fallen back into the corresponding solutions after a few iterations. This situation changes as we approach the boundary of the coexistence region. For smaller U , for example, the insulating solution becomes less and less attractive, until the basin of attraction vanishes at U_{c_1} . This scenario simply corresponds to the familiar free-energy landscape at a first-order transition.

Both the ED and the QMC discretize some component of the DMFT equations. In ED, the discretization parameter is the number n_s of *sites* of the quantum impurity model, while in QMC the number L of *slices* appears, with $\Delta\tau = \beta/L$ (cf. [2]). I have computed the insulating bath Green’s function $\mathcal{G}_0^{ins}(\tau)$ at the reference point for $n_s = 5, 6, 7$. Each of the solutions (for fixed n_s) is fully converged, in addition, the convergence with the number of sites n_s is excellent. I stress in passing that the sites in the ED algorithm are chosen in an optimal way, very similar to what is done in Gaussian integration [8]. Exponential convergence in n_s has been reported [2]. Given that $\mathcal{G}_0^{ins}(\tau, n_s = 7)$ differs by less than 0.005 from $\mathcal{G}_0(\tau, n_s = 6)$, I am led to the proposition that both $\mathcal{G}_0^{ins}(\tau, n_s = 7)$ and $\mathcal{G}_0^{met}(\tau, n_s = 7)$ are essentially exact. This assertion would have to be refuted for at least one

*krauth@physique.ens.fr, <http://www.lps.ens.fr/~krauth>

of the solutions (in fact, for the insulating one) if we were to agree with the authors of ref. [3].

The following reasoning will lead us to a very interesting result: consider a *single* iteration of the self-consistency loop

$$\mathcal{G}_0^{ins,met} \longrightarrow G \longrightarrow \mathcal{G}_0^{\text{new}} \quad (1)$$

If, contrary to our assertion, $\mathcal{G}_0^{ins}(\tau)$ or $\mathcal{G}_0^{met}(\tau)$ were not self-consistent, we should be able to detect differences, say, between $\mathcal{G}_0^{ins}$ and $\mathcal{G}_0^{\text{new}}$ - even by other methods than ED, for example by QMC simulation.

In this sense, I have discretized both $\mathcal{G}_0^{ins}$ and $\mathcal{G}_0^{met}$ and used them as inputs for single self-consistency loops (as in eq. (1)) of the QMC algorithm for different values of $\Delta\tau$. From the discretized Green's functions $G^{\Delta\tau}(\tau)$, we then compute $\mathcal{G}_0^{\text{new}}(\tau)$ by inverse Fourier transformation. Results for the metallic solution are shown in Fig. 1. We see that the finite- $\Delta\tau$ effects are quite small. Clearly, $\mathcal{G}_0^{\text{new}}(\tau, \Delta\tau) \rightarrow \mathcal{G}_0^{met}(\tau)$ as $\Delta\tau \rightarrow 0$. The metallic solution has not been contested.




FIG. 1. $\mathcal{G}_0^{\text{new}}(\tau, \Delta\tau)$ of the metallic solution, as computed by QMC for $\Delta\tau D/\sqrt{2} = 1, 0.5$ and $\mathcal{G}_0^{met}(\tau, n_s = 7)$ (from below). Parameters of the reference point are $\beta D/\sqrt{2} = 100, U = 3.55D/\sqrt{2}$.

The situation becomes much more illuminating as we consider the insulating solution $\mathcal{G}_0^{ins}(\tau)$. In this case, I have computed $G^{\Delta\tau}(\tau)$ for $\Delta\tau = 1, 0.5, 0.25$. The results for $\mathcal{G}_0^{\text{new}}(\tau)$, again obtained by inverse Fourier transformation, are shown in Fig. 2: it is evident that very large finite- $\Delta\tau$ effects are present. Notice that *all* the curves $\mathcal{G}_0^{\text{new}}(\tau, \Delta\tau)$ expose a plateau $\mathcal{G}_0 \sim \text{constant}$ for τ away from 0 or β . Such a plateau is characteristic of an insulating solution (cf. Fig. 1), and its value *decreases* with the gap of the single-particle density of states. We thus arrive at the crucial observation that the finite- $\Delta\tau$ effects bias the self-consistency condition of the QMC algorithm towards the metallic solution. Nevertheless, we find back the ED solution as $\Delta\tau \rightarrow 0$.




FIG. 2. $\mathcal{G}_0^{\text{new}}(\tau, \Delta\tau)$ of the insulating solution, as computed by QMC for $\Delta\tau D/\sqrt{2} = 1, 0.5, 0.25$ and $\mathcal{G}_0^{iso}(\tau, n_s = 7)$ (from below) at the reference point.

In terms of the free-energy landscape mentioned earlier, this simply means that the picture at finite $\Delta\tau$ is *tilted*. The QMC iteration is dragged away from the insulating solution $\mathcal{G}_0^{ins}$ into the metal.

Besides this effect, there seems to be no difference between the iteration loop of ED and the one of QMC. Within a deep basin of attraction of the insulating solution, a small drag due to finite $\Delta\tau$ effects should only lead to a shift of the solution, and its stability should be preserved. This is exactly what I have observed. I have run *full* iteration loops starting from $\mathcal{G}_0^{ins}(\tau)$ for $\Delta\tau D/\sqrt{2} = 1$ and $\Delta\tau D/\sqrt{2} = 0.5$. In the first case, the simulation moved away from the insulation solution: After about 10 iterations, the metallic solution was approximately recovered. In contrast, for $\Delta\tau D/\sqrt{2} = 0.5$, the insulating solution is very clearly stable. A 40-day simulation of this single problem on a work station has obtained a very well-converged self-consistent solution for $\mathcal{G}_0(\tau, \Delta\tau)$ within QMC. This solution makes no more reference to ED, but it of course resembles the curves shown in Fig. 2; the plateau value is $\mathcal{G}_0(\tau D/\sqrt{2} = 50) \sim 0.30$, comparable to the one-shot solution at the same value of $\Delta\tau$.

In my opinion, the present discussion of the discretization errors and both the one-step iteration and the explicit self-consistent Monte Carlo solution leave little room but to accept the coexistence at finite temperature. It is evident that the Monte Carlo simulation at the insulating solution has important finite- $\Delta\tau$ effects. These errors modify the qualitative aspects of the solution only close to the phase boundaries where the basins of attraction of the insulating solution are shallow, and small. Therefore, the discretization errors will be more pronounced close to U_{c1} , and also at higher temperature. This is what happened in the simulation of ref. [3], and what led the authors to wrong conclusions.

Acknowledgement: it is a pleasure to thank G. Kotliar and A. Georges for helpful discussions.

- [1] The correct scaling of energies was proposed in: W. Metzner and D. Vollhardt, *Phys. Rev. Lett.* **62**, 324 (1989); the self-consistent theory was formulated in: A. Georges and G. Kotliar, *Phys. Rev. B* **45**, 6479 (1992); solutions of the equations were first obtained in: M. Jarrell, *Phys. Rev. Lett.* **69**, 168 (1992); M. J. Rozenberg, X. Y. Zhang, and G. Kotliar, *Phys. Rev. Lett.* **69**, 1236 (1992); A. Georges and W. Krauth, *Phys. Rev. Lett.* **69**, 1240 (1992).
- [2] A. Georges, G. Kotliar, W. Krauth, and M. J. Rozenberg, *Rev. Mod. Phys.* **68**, 13 (1996).
- [3] J. Schlipf, M. Jarrell, P. G. J. van Dongen, N. Blumer, S. Kehrein, Th. Pruschke, and D. Vollhardt, *Phys. Rev. Lett.* **82** 4890 (1999).
- [4] M. Caffarel and W. Krauth, *Phys. Rev. Lett.* **72** 1545 (1994).
- [5] M. J. Rozenberg, R. Chitra, G. Kotliar, cond-mat/9906131.
- [6] L. Laloux, A. Georges and W. Krauth, *Phys. Rev. B* **50**, 3092, (1994).
- [7] The solutions $\mathcal{G}_0(\tau)$ are made available on request.
- [8] W. H. Press, S. A. Teukolsky, W. T. Vetterling, and B. P. Flannery, *Numerical Recipes*, 2nd edition, Cambridge University Press (1992).